

RELATIONSHIP BETWEEN PATIENT'S SATISFACTION AND LEVEL OF EMOTIONAL INTELLIGENCE IN NURSES WORKING IN TERTIARY CARE HOSPITAL

LAHORE

Orung Zaib Masih, Madiha Mukhtar, Sarfraz Masih

MSN Student at Lahore School of Nursing, The University of Lahore

Assistant Professor at Lahore School of Nursing, The University of Lahore

Principal at Lahore School of Nursing, The University of Lahore

ABSTRACT

Background: Emotional intelligence is a social skill and an important component of communication. It can be defined as the ability to correctly recognize one's own and others' emotions, as well as the ability to understand, evaluate, and use that information to make better decisions, solve problems, and live a better life. Nursing is a career that necessitates a high level of critical thinking and intellect, as well as a combination of relative intelligence and emotional intelligence. Emotional intelligence is the ability to sense what is going on, to understand how it feels while in that circumstance, to effectively analyse and consider other people's thoughts and feelings, and to respond accordingly.

Objectives: To determine the association between patient's satisfaction and level of emotional intelligence (EI) in nurses working in medical and surgical department in tertiary care hospitals, Lahore

Methodology: A correlational study design was used to carry on this study. This study was conducted at Mayo Hospital Lahore. The study participants were registered nurses (both male and female) and admit patients (both male and female) there. Purposive sampling technique was used n=265 participants were recruited. To collect data, first of all, permission was granted from the Research Ethical Committee (REC) of the University of Lahore. Then permission was taken

from the Medical superintendent of Mayo Hospital Lahore. To assess the performance among nurses and patients, an observational checklist was used. Data of the study participants was entered in statistical software SPSS 21 and was analyzed accordingly.

Results: This study shows that there are statistically non-significant differences between patient's satisfaction and emotional intelligence. The results revealed a non-significant association between nurses' emotional intelligence and patients satisfaction, as P -value is $P=.381$ which is non-significant. So, we reject the alternative hypothesis and accept the null hypothesis.

Conclusion: It is concluded that nurse's emotional intelligence has no relationship on patient's satisfaction. Nurse's emotional intelligence and its association with patient's satisfaction, which results were non-significant.

Key words: Emotional Intelligence, Patient's Satisfaction, Nurses

INTRODUCTION

Emotional intelligence is a type of cognitive ability that combines personality traits and social abilities to help people comprehend others' feelings and make better personal and organizational decisions. People with high emotional intelligence are better able to deal with organizational issues (Uhrich et al., 2021). Emotional intelligence describes how people deal with positive and negative life situations, as well as how they carry out efficient and long-term actions in the workplace (Magnano et al., 2016).

Emotional intelligence is seen as a necessary component of nursing practice in the United States (Hamid et al., 2015). Emotional intelligence is defined as the ability of people to recognize, understand and control their own emotions and recognize, understand and influence the emotions of others (Al-Ruzzieh & Ayaad, 2021). Emotional Intelligence (EI) is an important skill for both personal and professional success. Self-awareness, self-management, social awareness, and relationship management are all characteristics of EI (Hadiwijaya & Hutasoit, 2017).

Nursing is a career that necessitates a high level of critical thinking and intellect, as well as a combination of relative intelligence and emotional intelligence (Faravani & Atai, 2015). As a result, in order to cope with the emotional demands of the healthcare setting, nurses must have high levels of emotional intelligence skills. Emotional intelligence develops the ability to deal with interpersonal issues by managing emotional responses, being empathic, and remaining attentive to patient concerns, resulting in more effective cooperation and better patient care (Foster & McCloughen, 2020).

Emotional intelligence can help nurses manage their own and their patients' emotions, demonstrate authentic emotional responses, are empathic and convey emotions without causing conflict, and manage instinctive emotions in nurse-patient encounters, such as disgust, anger, and frustration. Nurses can handle a variety of clinical scenarios by seeing things from the patients' perspective and empathizing with their feelings (Beauvais et al., 2017). Patients who are cared for by a nurse who exhibits EI skills believe that the nurse genuinely cares about their well-being and health, which is the essence of nursing and caring (Celik, 2017).

Patient satisfaction is a measurable criterion for assessing health care and, as a result, nursing care quality (Karaca & Durna, 2019). Patient-centred care is the most important factor of care quality (PCC). Patient-centered care involves six critical components, according to the Institute of Medicine: safety, efficacy, time, efficiency, quality, and equity. The conduct and emotions of nurses have a direct impact on the quality of patient care (Richards & Goldin, 2015).

In recent years, many studies have been conducted to determine the level of patient satisfaction with nursing care. For instance, studies done in Iraq (M Mussa et al., 2016), Brazil and Egypt showed that patient satisfaction with nursing care was high. Additionally, the overall level of patient satisfaction with nursing care was 69% in Iran, 67% in Kenya, and 33% in Ghana (Mulugeta et al., 2019). On the contrary, the results of the study done in India revealed that most of the hospitalized patients had poor perception regarding nursing care (Rajalatchumi et al., 2018).

MATERIAL AND METHODS

A Correlational study design was used to carry on this research work. This study was conducted at the Mayo Hospital Lahore and Social Security hospital Lahore. The study participants were all the Nurses (male and female) and admit patients from medical and surgical departments. A purposive sampling technique was used to recruit the recommended sample. Sample size of 133 nurses and 265 patients has been calculated with 95% confidence interval.

Inclusion Criteria: Nurses (male and female) who are working in medical and surgical department. Age 24-60 years of Nurses .Have one year of work experience .Patients included the following: a minimum of 48 hours stay time in the hospital .Patients; aptitude for reading and responding to the questions .Only medical department patients. Patient's Full alertness (aware of person, place, and time); no history of psychiatric illnesses. Exclusion Criteria Nursing Students Nursing administrative. Loss of consciousness, abrupt illness progression, cardiac or respiratory arrest, and a wish to stop the study. Other department's nurses and patients. Tool 1. Emotional Intelligence (EI) of Nurses. It comprises on 33-item self-reports that is brief EI. The scale's responses, which were of the five-point Likert variety, were Strongly Agree, Agree, Neither, Disagree, and Strongly Disagree (Tatar et al., 2011). Below 75 EI Score = <45% was consider Low EI. Above 75-125 EI Score = >50% -78% was consider Moderate EI. Above 125-165 EI Score = >78% -100% was consider High EI. Tool 2. Patients Satisfaction. It comprises on 18-item self-reports that is brief patients satisfaction. The scale's responses, which were of the five-point Likert variety, not satisfied, rarely satisfied , satisfied, extremely satisfied , and fully satisfied (Akin & Erdogan, 2007). Below 45 Score = < 50 % was consider Low Satisfaction, Above 45-70 Score = > 50% -78% was consider Moderate Satisfaction, Above 70-90 Score = >78% -100% was consider High Satisfaction. To collect data, first of all permission was granted from the Research Ethical Committee (REC) of the university of Lahore. Then permission was taken from Medical superintendent of Mayo Hospital Lahore and social security hospital Lahore. Data of the study participants were entered in statistical software SPSS 21.0 and was analysed accordingly. Result of the study were presented as median through table. Parsons correlation

coefficient was applied to checked correlation. P-value ≤ 0.05 will be considered statistically significant.

RESULT

Table: 1 below indicated that there were 133 participants were involved in the study. The majority, 42.9%, of participants were between the ages of 24-30 years, 39.8% of participants were from the age group of 31-40, and 12.8% participants were from the age group of 41-50, while 6 % participants were from the age group of 51-60. Most participants were female, with (n=94) 70.3% of the population. Male participants were (n=39) with 29.3 % of the population. Married, with (n=84) 63.2% of the population. Unmarried participants were (n=49) with 36.8 % of the population. Revealed that the majority, 50.4%, of participants were diploma holders, 49.6% were Post RN BSN. The results of total years of job experience. Results revealed that 30.41% of participants had an experience of <Then 5 Years. Most 36.8 % of participants had experience of 6-10 years, whereas 16.5% had an experience of > 11-15 and 11.3% participant had an experience >16-20 1 year also 7 % participant experience more the > 20 years. Most of the 66.2% of participants had belonged medical department, while 33.8% had belonged surgical department.

Table:1 Socio-Demographic Characteristics of Nurse's

Variables	Frequency
Age of nurses in years	(n=133)
24-30 years	57(42.9%)
31-40 years	53 (39.8%)
41-50 years	12 (12.8%)
51-60 years	6 (4.6%)
Gender	(n=133)
Male	39 (29.3%)
Female	94 (70.7%)
Marital Status	(n= 133)
Married	84 (63.2%)
Unmarried	49 (36.8%)
Qualification	(n= 133)
Post RN BScN	66 (49.4%)
Diploma in Nursing	67 (50.6%)
Job experiences	(n= 133)

1-5 Years	40 (30.1%)
6-10 Years	49 (36.8%)
11- 15 Years	22 (16.5%)
16- 20 Years	15 (11.3%)
Above 20 Years	7 (5.3%)
Departments	(n= 133)
Medical	88(66.2%)
Surgical	45 (33.8%)

Table 2: Below demographic Characteristics of patient's participants. 265 participants were involved in this study. The majority, 44.5%, of participants were between the ages of 30-40 years, 28.3% of participants were from the age group of 18-30, 20.4% participants were from the age group of 40-50, while 6.8 % participants were from the age group of 51-60. Most participants were female, with (n=145) 54.7% of the population. Male participants were (n=120) with 45.3 % of the population. Revealed that the majority, 56.4%, of participants were high school, 24.9% were middle school and 18.5 % were primary school.

Table:2 Socio-Demographic Characteristics of Patient's

Variables	Frequency
Age of Patients in years	(n=265)
18-30 years	75(28.3%)
31-40 years	118 (45.5%)
41-50 years	54(20.4%)
51-60 years	18(6.8%)
Gender	(n=265)
Male	120(45.3%)
Female	145(54.7%)
Qualification	(n= 265)
Primary School	49 (18.9%)
Middle School	66 (24.9%)
High School	150(56.6%)

Below table 3 indicated that 0 % nurses have EI and 72.9% of nurses have moderate EI, 27.1. % of participants has high EI.

Table: 3 Level of emotional intelligence in nurses

Variables	Frequency
EI of nurses	(n=133)
Low EI (< 45)	0 (0.0%)
Moderate EI (46-76 %)	97(72.9%)
High EI (76-100 %)	36 (27.1%)

Table 4 below shows that 15.1% patients have low satisfaction, 84.9 % of participants have moderate satisfaction and 0% patients have high satisfaction.

Table: 4 Level of Patients Satisfaction

Variables	Frequency
Patients Satisfaction	(n=265)
Low Satisfaction (< 50%)	40 (15.1%)
Moderate Satisfaction (< 50-70%)	225(84.9%)
High Satisfaction (>70-90%)	0 (0.0%)

Table 5 indicated that nurse's emotional intelligence and its association with patient's satisfaction, which results were non-significant as the p-value was .381 (P value is >0.05 taken as non-significant). There is no association regarding patient's satisfaction and level of emotional intelligence (EI) in nurses. The actual outcome of the Pearson's Correlation test is presented in below table. The results revealed a non-significant association between nurses' emotional intelligence and patients satisfaction, as P-value is P=.381 which is non-significant. So, we reject the alternative hypothesis and accept the null hypothesis.

Table: 5 Association among Nurses' Emotional Intelligence and Patients Satisfaction (n=398)

Correlations			
		Total EI	Total PS
Total EI	Pearson Correlation	1 133	.076 .381 133
	Sig. (2-tailed)		

	N		
Total PS	Pearson	.076	1
	Correlation	.381	265
	Sig. (2-tailed)	133	
	N		

DISCUSSION

The finding of the current study revealed that the nurse's age ranged between 24 to 60 years old. The majority (44.5%) of the nurses in the sample had their ages ranged between 30 and 40 years old. The majority of the participants were female. These results were similar to the results of Han, H., & Johnson, S. D. (2012) who studied the relationship. Their study showed that the participant ages ranged from 24 to 63 ($M=40.5$ $SD=10$). The participants represent the population in the program because the population was also primarily female (67.5%) (Han, 2009).

Atilla et al, in their study, showed that there was a positive and close relationship, with 0.01 significance level, between the self-evaluation and regulation of emotions sub dimensions of EI and the nursing satisfaction sub dimension of patient satisfaction. These findings are also consistent with our study results ($P=0.00$) (Atilla et al., 2013).

In a study, Karakas and Kucukogly measured the EI of nurses, and they reported that the nurses who took part in their study had low scores in the sub dimensions of controlling emotions and social skills and that those dimensions needed to be developed. In the present study, the nurses had values above the mean scores in the sub dimensions of EI (Cronbach's $\alpha =0.93$, mean EI: 3.96 ± 0.45) (KÜÇÜKOĞLU, 2011).

CONCLUSION

In conclusion, our study results show that as the level of EI nurses almost moderate, the satisfaction level of patients also moderate for whom the nursing care is provided also increases. Thus, patient satisfaction is affected by the EI level of the health care providers. Furthermore, it should be kept in mind that the EI skills are not endowments and that they can be learned and developed later. Therefore,

EI should be one of the determinants of the objectives and recognized as one of the quality indicators to improve the quality of health care services. The nurses should be given education periodically in order to improve and use EI.

REFERENCES

- Uhrich, B. B., Heggstad, E. D., & Shanock, L. R. (2021). Smarts or trait emotional intelligence? The role of trait emotional intelligence in enhancing the relationship between cognitive ability and performance. *The Psychologist-Manager Journal*, 24(1), 23.
- Al-Ruzzieh, M. A., & Ayaad, O. (2021). Impact of nurses' emotional intelligence on the implementation of a professional practice model in cancer care. *British Journal of Nursing*, 30(19), 1110-1116.
- Hadiwijaya, H., & Hutasoit, G. (2017). *Effect of emotional intelligence on student learning achievement*. Muhammadiyah University Metro.
- Faravani, A., & Atai, M. R. (2015). Multiple intelligences, dialogic-based portfolio assessment, and the enhancement of higher order thinking. *Journal of Teaching Language Skills*, 33(4), 19-44.
- Magnano, P., Craparo, G., & Paolillo, A. (2016). Resilience and Emotional Intelligence: which role in achievement motivation. *International Journal of Psychological Research*, 9(1), 9-20.
- Hamid, A., Hajmohammadi, S., Mehri, L., & Naddaf, S. (2015). The investigation between Emotional intelligence and job stress and quality of life among high school teachers in Khozestan province. *European Online Journal of Natural and Social Sciences: Proceedings*, 4(1 (s)), pp. 1371-1379.
- Foster, K. N., & McCloughen, A. J. (2020). Emotionally intelligent strategies students use to manage challenging interactions with patients and families: A qualitative inquiry. *Nurse Education in Practice*, 43, 102743.

- Beauvais, A., Andreychik, M., & Henkel, L. A. (2017). The role of emotional intelligence and empathy in compassionate nursing care. *Mindfulness & Compassion*, 2(2), 92-100.
- Celik, G. O. (2017). The relationship between patient satisfaction and emotional intelligence skills of nurses working in surgical clinics. *Patient preference and adherence*, 11, 1363.
- Cheshire, M. H., Strickland, H. P., & Ewell, P. J. (2020). Measured emotional intelligence in Baccalaureate Nursing Education: A longitudinal study. *Nursing Education Perspectives*, 41(2), 103-105.
- Karaca, A., & Durna, Z. (2019). Patient satisfaction with the quality of nursing care. *Nursing open*, 6(2), 535-545.
- M Mussa, Y., A Mohammed, H., & I Ali, R. J. K. U. J.-S. S. (2016). Assessment of adult patients satisfaction regarding nursing Care in Different Hospitals in Kirkuk City. *11(3)*, 222-236.
- Mulugeta, H., Wagnew, F., Dessie, G., Biresaw, H., & Habtewold, T. D. J. B. n. (2019). Patient satisfaction with nursing care in Ethiopia: a systematic review and meta-analysis. *18(1)*, 1-12.
- Rajalatchumi, A., Ravikumar, T. S., Muruganandham, K., Thulasingham, M., Selvaraj, K., Reddy, M. M., . . . medicine. (2018). Perception of patient safety culture among health-care providers in a tertiary care hospital, South India. *9(1)*, 14.
- Han, H. Y. (2009). *Relationship between students' emotional intelligence, social bond, and interactions in online learning*. University of Illinois at Urbana-Champaign.
- Atilla, G., ÇARIKÇI, İ. H., & Erdem, R. J. A. K. Ü. İ. V. İ. B. F. D. (2013). Hastanelerde duygusal zekâ-hasta memnuniyeti ilişkisi: Isparta il merkezi örneği. *15(1)*, 101-119.
- KÜÇÜKOĞLU, S. J. A. H. v. S. B. D. (2011). Bir eğitim hastanesinde çalışan hemşirelerin duygusal zeka düzeyleri. *14(3)*, 8-13.
- Richards, M. K., & Goldin, A. B. (2015). Patient-centered care and quality: Activating the system and the patient. *Seminars in pediatric surgery*,