

BIODIVERSITY OF ORDER COLEOPTERA AT SELECTED LOCALITIES OF DISTRICT BATTAGRAM THROUGH MALAISE TRAPPING SYSTEM

Syed Sabir Hussain Shah*, Asad Khan**, Hidayat Ullah*, Ateeq Ullah*, Irfan Khan*, Sana Ullah*, Izhar Ullah*, Muhammad Idrees*, Gurmeet Singh*, Fazal Amin*, Sajidur Rahman*, Bilal Ahmad*

*Department of Zoology Hazara University Mansehra Khyber Pakhtunkhwa, Pakistan

** Department of Zoology, Shaheed Benazir Bhutto University (SBBU) Sheringal, Dir Upper, Khyber Pakhtunkhwa, Pakistan

Abstract- The most populous taxon in the animal kingdom are insects. As the largest animal category, insects are collected using a number of sampling techniques. Among several methods of sampling for the collection of various insect groups, sweep nets, light traps, pitfall traps, Winkler sampling, and malaise traps are common and very effective. Coleoptera is the largest order of class insecta and have more than 400,000 known species. The most popular nonattractant, static insect traps are Malaise trap, which are big, tent-like structure made of fine mesh netting. District Battagram were selected for the installation of Malaise trap. Malaise trap were installed in five different sites (Biland kot, Kuza banda, Meran, Pirhari, Ajmera) for five months (4 April 2022 to 29 August 2022). Total 192 samples were trapped of Order Coleoptera which were further identified to seven families and to 12 genera. Out of these families, the family Scarabaeidae were founded with the highest number of specimens and the family Carabidae, family Chrysomelidae and family Coccinellidae were founded with the lowest number of specimens.

Index Terms- Animal, Coleoptera, Diversity, Insects, Malaise trap,

I. INTRODUCTION

The most populous taxon in the animal kingdom are insects [1]. As the largest animal category, insects are collected using a number of sampling techniques. Among several methods of sampling for the collection of various insect groups, sweep nets, light traps, pitfall traps, Winkler sampling, and malaise traps are common and very effective [2]. There are around 7,51000 recognized species of insects on the universe, which is almost one-fourth of all animal species [3]. The Coleoptera, which comprises 3,87,100 species and represents for about 38% of all insect orders, which is the most effective one [4].

ORDER COLEOPTERA

Coleoptera is the largest order of class insecta and have more than 400,000 known species [5]. This order has a great deal of diversity in terms of size, shape, biology, and behavior. They have two types of wings, a pair of fore wings also called elytra

and a pair of membranous hind wings. They have complete metamorphosis and have chewing mouthparts which are modified for various functions. Coleoptera are found in various habitat like desert, water and feed on various plant and animal materials [6].

MALAISE TRAP

The most popular nonattractant, static insect traps are Malaise trap, which are big, tent-like structure made of fine mesh netting. Rene Malaise invented them when he observed insects in a tent that provided the first inactive method for observing and gathering a large number of flies. Several bug species will climb up when they encounter a barrier, which is advantageous for Malaise trap. The mesh walls of the traps stop flying insects, particularly Diptera and Hymenoptera, although they also catch a variety of ground-dwelling species [7]. Malaise traps can be set up everywhere there are likely to be flying insects, such as woodlands, open grasslands, prairies, and windy summits, but care must be taken to secure them in the event of strong winds. The placement of the traps has an impact on the biodiversity collection [8]. The traps can be used for general gathering, biodiversity surveys, and research into the migration of insects, and both nocturnal and diurnal patterns of abundance, as well as vertical stratification. The species that are gathered are affected by numerous variables. Abiotic variables like the weather and season can have a variety of consequences on the behavior of the target taxa and the species that are trapped. The species that are trapped will depend on the design, placement, and orientation of the trap as well as the addition of various lures. The kind and amount of plants nearby as well as species-specific behavior are biological factors that affect trap catch. Taxon-specific factors determine whether Malaise traps are superior or inferior to other trap types in terms of their ability to trap specific groups. Because they have the least taxon overlap with Malaise traps, pitfall traps and leaf litter extraction are good choices to couple with Malaise traps to enhance the amount of arthropod diversity tested while putting out the least amount of effort. Yet, there are

certain drawbacks to using malaise traps. Commercially available traps cost a lot of money and are easily damaged by large mammals like bears or by high winds. Also, because Malaise traps frequently gather enormous quantities of arthropods, there may be concerns about the trapped and demise of non-target by catch [9].

II MATERIALS AND METHODS

STUDY SITE

District Battagram were selected for the installation of Malaise trap. Battagram is situated in the western part of Khyber Pakhtunkhwa, Pakistan, with a total surface area of 1301 square kilometres, between 34° 33 and 34° 47' N and 72° 54 and 73° 15' E [10]. Mansehra district, Kohistan district, district of Torghar, and Shangla district are its borders on the east, north, south, and west, respectively. The Battagram district has a population of 476,612 as per the 2017 Census. Green, luscious mountains may be found at Battagram. At Thakot and Sukaisar, the altitude is above 4690 m, respectively, with an altitude of 525 m. A variety of climatic conditions exist, ranging from subtropical to "alpine" climates. There are many agricultural fields, wastelands, forests, and meadows in Battagram [10]. Agriculture is the most common form of living for most people, which is followed by the farming of animals. They grow wheat, red beans, vegetables, rice, corn and rice [11].

MALAISE TRAP DESIGN AND SETUP

Four wooden rods were buried in the ground in a square shape at a distance of 4 feet from each other. All the rods were covered with black cloth by three sides and one side was left opened. Black cloth attracts insects more than other cloths, so it is important to use black cloth in the trap. While the upper part of the trap were covered with white net like cloth. The four rods connect with each other by four wooden rods at the upper portion. A wooden rod turned into a cone structure and a bottle was attached to the cone and this bottle contained ethanol. Insects used to fly on the trap. Insects kept falling in the ethanol filled bottle when they move upward. All the insects were collected through this trap.



SAMPLING, PRESERVATION AND IDENTIFICATION

Forceps were used to remove all of the insect samples from the Malaise trap, and they were stored in small bottles. The trapped specimens were kept in 70% ethanol in small bottles. The trapped samples were kept in 70% ethanol in small bottles. The same species of insects were preserved in same one bottle. All the bottles were labelled with Order name, date and locations. The bottles were labelled with order name, date and locality. The specimens were further stored in fridge at 21°C in the Zoological Laboratory of Hazara University Mansehra. The trapped samples were first identified with the help of microscope. Up to Order levels in the Zoological Museum of Hazara University Mansehra. Up to species level, the specimens were identified in National Insect Museum (NIM) in National Agriculture Research Center (NARC) Islamabad.


III RESULTS

A total of 192 specimens of Order Coleoptera were trapped in five months (4 April 2022 to 29 August 2022) which were identified in National Insect Museum, National Agricultural Research Center Islamabad, Pakistan. 192 Specimens of Order Coleoptera belonging to seven different families which were family Carabidae, family Cerambycidae, family Chrysomelidae, family Coccinellidae, family Hybosoridae, family Meloidae and family Scarabaeidae. The Malaise trap were installed in five different localities (Biland kot, Kuza banda, Meran, Pirhari, Ajmera) of District Battagram.

FAMILY WISE RICHNESS

192 specimens were belonging to seven different families of order Coleoptera. Out of seven families, the family Scarabaeidae have highest number of specimens and the family Carabidae,


family Chrysomelindae and family Coccinellidae have the lowest number of specimens.



Family wise richness


GENUS WISE RICHNESS

The total specimens of seven families were further identified to 12 different Genera. Out of 12 Genera, the Genus Mylabris have the highest number of specimens and the Genus Pheropsophus have the lowest number of specimens.



LOCALITY WISE RICHNESS

The maliase trap were installed in five different localities of District Battagram. Out of these localities, maximum specimens were trapped at Biland kot and manimum specimens were trapped at Ajmera. The locality wise richness are given below.



IV ACKNOWLEDGMENT

AUTHORS ACKNOWLEDGMENT: DEPARTMENT OF ZOOLOGY HAZARA UNIVERSITY MANSEHRA KHYBER PAKHTUNKHWA, PAKISTAN

V REFERENCES

- [1] E. O. Wilson, "The little things that run the world (the importance and conservation of invertebrates)," ed: JSTOR, 1987, pp. 344-346.
- [2] S. E. Fraser, C. Dytham, and P. J. Mayhew, "Determinants of parasitoid abundance and diversity in woodland habitats," *Journal of applied ecology*, vol. 44, no. 2, pp. 352-361, 2007.
- [3] A. Choudhary and J. Ahi, "Biodiversity of freshwater insects: a review," *The International Journal of Engineering and Science*, vol. 4, no. 10, pp. 25-31, 2015.
- [4] Z.-Q. Zhang, *Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness*. Magnolia press, 2011.
- [5] P. Bouchard *et al.*, "Family-group names in Coleoptera (Insecta)," *ZooKeys*, no. 88, p. 1, 2011.
- [6] P. J. Gullan and P. S. Cranston, *The insects: an outline of entomology*. John Wiley & Sons, 2014.
- [7] K. van Achterberg, "Can Townes type Malaise traps be improved? Some recent developments," *Entomologische berichten*, vol. 69, no. 4, pp. 129-135, 2009.
- [8] S. R. Leather, *Insect sampling in forest ecosystems*. John Wiley & Sons, 2008.
- [9] M. J. Skvarla, J. L. Larson, J. R. Fisher, and A. P. Dowling, "A review of terrestrial and canopy Malaise traps," *Annals of the Entomological Society of America*, vol. 114, no. 1, pp. 27-47, 2021.

- [10] F. Haq, "The critically endangered flora and fauna of district Battagram Pakistan," *Advances in Life sciences*, vol. 2, no. 4, pp. 118-123, 2012.
- [11] S. Muhammad, "Resource Management Plan Allai forests," 2004.

AUTHORS

First Author – Syed Sabir Hussain Shah, Master of Philosophy in Zoology, Hazara University Mansehra, Pakistan

Second Author – Asad Khan, Department of Zoology, Shaheed Benazir Bhutto University (SBBU) Sheringal, Dir Upper, Khyber Pakhtunkhwa, Pakistan

Third Author – Hidayat Ullah , Doctor of Philosophy in Zoology, Hazara University Mansehra, Pakistan

Correspondence Author – Syed Sabir Hussain Shah,